

# A model for conservative chaos constructed from multi-component Bose-Einstein condensates with a trap in 2 dimensions

Hisatsugu Yamasaki\* and Katsuhiro Nakamura  
*Department of Applied Physics, Osaka City University  
Sumiyoshi-ku Osaka, 558-8585 Japan*

Yuhei Natsume  
*SGraduate School of Science and Technology, Chiba University  
Inage-ku, Chiba, 263-8522 Japan*

(Dated: February 2, 2008)

To show a mechanism leading to the breakdown of a particle picture for the multi-component Bose-Einstein condensates (BECs) with a harmonic trap in high dimensions, we investigate the corresponding 2- $d$  nonlinear Schrödinger equation (Gross-Pitaevskii equation) with use of a modified variational principle. A molecule of two identical Gaussian wavepackets has two degrees of freedom (DFs), the separation of center-of-masses and the wavepacket width. Without the inter-component interaction (ICI) these DFs show independent regular oscillations with the degenerate eigen-frequencies. The inclusion of ICI strongly mixes these DFs, generating a fat mode that breaks a particle picture, which however can be recovered by introducing a time-periodic ICI with zero average. In case of the molecule of three wavepackets for a three-component BEC, the increase of amplitude of ICI yields a transition from regular to chaotic oscillations in the wavepacket breathing.

Recently a great number of theoretical and experimental efforts have been devoted to Bose-Einstein condensates (BECs) [1, 2]. As well as single-component BECs, the trapping techniques can create multi-component condensates which involve inter-component nonlinear interactions. The multi-component BEC, far from being a trivial extension of the single-component one, presents novel and fundamentally different scenarios for its ground state and excitations. In particular, it has been observed that BEC can reach an equilibrium state characterized by separation of the components in different domains [3].

BEC has a dual aspect of waves and particles. The wave nature is high-lighted in the phenomenon of interference leading to fringe patterns [1, 2]. On the other hand, the particle nature of BECs can be seen in typical localized states like vortices and solitons. In fact solitons were observed in the quasi-one dimensional BEC [4, 5].

Among the works that emphasize a role of the particle picture for BEC *with a harmonic trap* in high-dimensions, those of Pérez-García *et al.* are the most noteworthy [6, 7, 8, 9]. We focus on their two important assertions. The first one made for a single-component BEC is as follows [8]: If the phase of BEC wave function will be suitably corrected, the center-of-mass for a wavepacket displaced from the origin obeys Newtonian dynamics and the Ehrenfest theorem is valid even for the nonlinear Schrödinger equation (NSE). Furthermore the center-of-mass is decoupled from dynamics of the shape of a wavepacket. The second assertion is concerned with the multi-component BEC [9]: If the distance between

wavepackets with each *linked to the individual component* is much larger than their typical widths, the picture of soliton molecules composed of solitonic atoms (wavepackets) holds for the multi-component BEC with a harmonic trap in high-dimensions. The large distance as above is guaranteed by a sufficiently large centrifugal force due to non-vanishing angular momentum. .

Pérez-García *et al.*'s second assertion is quite interesting because the solitonic structures like wavepackets are believed to be dynamically unstable in two and higher spatial dimensions. However, we have several criticisms against their assertion: (1) Inter-component interaction has a tendency to swell out individual wavepackets and breaks their picture of interacting particles, as evidenced in Fig.1(a); (2) Collective coordinates for the width and phase of wave packets, which should be coupled with the center-of-mass, are not taken into consideration, although being studied intensively in other works of their own [6, 11].

In this Letter, choosing the 2- $d$  multi-component BEC with a harmonic trap, we develop a refined variational principle to derive an effective dynamics for interacting wave packets and pinpoint the weak points of the idea of Pérez-García *et al.* [9]. Then we shall propose a new model to restore the picture of soliton molecules in two dimensions. In particular, by controlling a parameter we shall see a transition of the wavepacket breathing from regular to chaotic oscillations. We numerically analyze the subject on the basis of a three-component BEC corresponding to the “three-body problem”.

The multi-component BEC at zero temperature is described by NSE (or Gross-Pitaevskii (GP) equation). We shall consider a system of  $n$  complex fields

---

\*Electronic address: hisa@a-phys.eng.osaka-cu.ac.jp




FIG. 1: (a) Time evolution (up to  $t = 30[\omega^{-1}]$ ) of initial three Gaussian wavepackets in 3-component BEC with angular momentum  $M = 9$  and  $g = \Lambda g = 2\pi$  (computed by using Eq.(1)). See an outbreak of wave interference; (b) Fat and skinny normal modes (schematic).

$\psi_1(t, \mathbf{r}), \psi_2(t, \mathbf{r}), \dots, \psi_n(t, \mathbf{r})$  ruled by the equations

$$i\partial_t \psi_j(t, \mathbf{r}) = \left[ -\frac{1}{2}\Delta_{\mathbf{r}} + \frac{\mathbf{r}^2}{2} \right] \psi_j(t, \mathbf{r}) + \sum_{j,k} g_{jk} |\psi_k(t, \mathbf{r})|^2 \psi_j(t, \mathbf{r}) \quad (1)$$

for  $j, k = 1, \dots, n$  (in units of atomic mass  $m = 1$ , confining length  $\sqrt{\frac{\hbar}{m\omega}} = 1$  and oscillation period  $\omega^{-1} = 1$ ).  $\mathbf{r}^2/2 = (x^2 + y^2)/2$  stands for a harmonic trap and  $\sum_k g_{jk} |\psi_k(t, \mathbf{r})|^2$  is the nonlinear term. The coefficient  $g_{ij}$  (proportional to the scattering length scaled by confining length) is tunable by Feshbach resonance. We first consider a static case of repulsive nonlinearity,  $g_{ij} > 0$  and choose  $g_{ii} = g$  and  $g_{ij} = \Lambda g$  for  $i \neq j$ .

In the absence of the inter-component interaction (ICI), i.e., for  $\Lambda g = 0$ , each component has a solitonic state, namely Gaussian wavepacket or vortex solution with a movable center-of-mass. On the other hand, the non-vanishing ICI ( $\Lambda g \neq 0$ ) among solitons may yield soliton-soliton interactions. To reveal a crucial role of the breathing nature of above localized structures, we apply a refined variational principle and derive from (1) the evolution equation for the collective coordinates of wavepackets. The trial Gaussian wavepackets (WPs) are constructed from the circularly-symmetric ground state solution of (1) with ICI suppressed:

$$\begin{aligned} \psi_j(\mathbf{r}) &= \sqrt{\frac{1}{\pi w_j^2}} \exp \left[ -\frac{(\mathbf{r} - \mathbf{R}_j)^2}{2w_j^2} \right] \exp(i\Theta_j(\mathbf{r})), \\ \Theta_j(\mathbf{r}) &= \boldsymbol{\alpha}_j(\mathbf{r} - \mathbf{R}_j) + \frac{1}{2}(\mathbf{r} - \mathbf{R}_j)^T \boldsymbol{\beta}_j(\mathbf{r} - \mathbf{R}_j). \end{aligned} \quad (2)$$

(One may also choose excited states responsible for vortices.)  $\mathbf{R} = (X, Y)$  is the center-of-mass and  $w$  is width of the circular WP.

$$\boldsymbol{\alpha} = (\alpha^X, \alpha^Y) \quad (3)$$

and

$$\boldsymbol{\beta} = \begin{pmatrix} \beta^{XX} & \beta^{XY} \\ \beta^{YX} & \beta^{YY} \end{pmatrix}, \quad \beta^{XY} = \beta^{YX} \quad (4)$$

are respectively the first- and second-order coefficients of Taylor-expansion of the phase  $\Theta$  w.r.t.  $\mathbf{r} - \mathbf{R}$  (a trivial constant phase is suppressed).  $\mathbf{R}$ ,  $w$ ,  $\boldsymbol{\alpha}$  and  $\boldsymbol{\beta}$  constitute collective coordinates below.

The expansion of  $\Theta$  in Eq.(2) is the most natural, although the existing works [6, 7, 10, 11] employ an expansion w.r.t.  $\mathbf{r}$  rather than  $\mathbf{r} - \mathbf{R}$ . The advantage of our expansion is that  $\boldsymbol{\alpha}$  and  $\boldsymbol{\beta}$  turn out to have a transparent correspondence with the velocity of center-of-mass and the WP breathing, respectively (see (8) and (9) below).

According to the variational principle, the action function deriving Eq.(1) is obtained from Lagrangian density for field variables:

$$\begin{aligned} \mathcal{L} &= \sum_{j=1}^n \left[ \frac{i}{2} (\psi_j \dot{\psi}_j^* - \psi_j^* \dot{\psi}_j) + \frac{1}{2} |\nabla_{\mathbf{r}} \psi_j|^2 \right. \\ &\quad \left. + \frac{\mathbf{r}^2}{2} |\psi_j|^2 + \frac{g}{2} |\psi_j|^4 \right] + \sum_{k>j} \Lambda g |\psi_j|^2 |\psi_k|^2. \end{aligned} \quad (5)$$

In fact, the multi-component GP equation is obtained from Lagrange equation,  $\frac{\partial}{\partial t} \frac{\partial \mathcal{L}}{\partial \psi_j^*} - \frac{\partial \mathcal{L}}{\partial \psi_j^*} + \nabla \frac{\partial \mathcal{L}}{\partial \nabla \psi_j^*} = 0$ .

We now insert (2) into (5) and obtain Lagrangian  $L$  for the collective coordinates by integrating  $\mathcal{L}$  over space coordinates:

$$\begin{aligned} L &= \int \mathcal{L} d\mathbf{r} = \sum_{j=1}^n \left[ \left( \text{Tr} \dot{\boldsymbol{\beta}}_j + \text{Tr} (\boldsymbol{\beta}_j^T \boldsymbol{\beta}_j) \right) \frac{w_j^2}{4} + \frac{\boldsymbol{\alpha}_j^2}{2} \right. \\ &\quad \left. - \boldsymbol{\alpha}_j \dot{\mathbf{R}}_j + \frac{\mathbf{R}_j^2}{2} + \frac{w_j^2}{2} + \left( 1 + \frac{g}{2\pi} \right) \frac{1}{2w_j^2} \right] \\ &\quad + U(\{\mathbf{R}_j\}, \{w_j\}) \end{aligned} \quad (6)$$

with

$$U(\{\mathbf{R}_j\}, \{w_j\}) = \sum_{j>k} \frac{\Lambda g}{\pi(w_j^2 + w_k^2)} e^{-\frac{(\mathbf{R}_j - \mathbf{R}_k)^2}{w_j^2 + w_k^2}}. \quad (7)$$

Lagrange equations of motion for phase variables  $\boldsymbol{\alpha}$  and  $\boldsymbol{\beta}$  lead to

$$\boldsymbol{\alpha}_j = \dot{\mathbf{R}}_j \quad (8)$$

and

$$\beta_j^{XX} = \beta_j^{YY} = \frac{\dot{w}_j}{w_j}, \quad \beta_j^{XY} = \beta_j^{YX} = 0. \quad (9)$$

Noting that  $\boldsymbol{\alpha}$  and  $\boldsymbol{\beta}$  in (8) and (9) (and also all higher-order phase coefficients) *adiabatically follow*  $\mathbf{R}$

and  $w$ , we can rewrite Lagrange equations for the latter by eliminating the former: Equation of motion for  $\mathbf{R}$ ,  $\frac{d}{dt} \left( \frac{\partial L}{\partial \dot{\mathbf{R}}_j} \right) - \frac{\partial L}{\partial \mathbf{R}_j} = 0$ , becomes

$$\ddot{\mathbf{R}}_j + \mathbf{R}_j + \frac{\partial}{\partial \mathbf{R}_j} U(\{\mathbf{R}_j\}, \{w_j\}) = 0. \quad (10)$$

Similarly for  $w$  we have  $\frac{d}{dt} \left( \frac{\partial L}{\partial \dot{w}_j} \right) - \frac{\partial L}{\partial w_j} = 0$ , which is reduced to

$$\ddot{w}_j + w_j - \frac{1}{w_j^3} \left( 1 + \frac{g}{2\pi} \right) + \frac{\partial}{\partial w_j} U(\{\mathbf{R}_j\}, \{w_j\}) = 0. \quad (11)$$

A couple of Eqs. (10) and (11) indicate: Contrary to Pérez-García *et al.*'s theory that ignored the role of WP width[9], the width and center-of-mass are strongly correlated, leading to a breakdown of the picture of a solitonic molecule based on the center-of-masses alone. To examine the problem in detail, we consider the two-component BEC in two dimensions and explore the fate of a WP breathing against ICI ( $\Lambda g$ ). Effective Lagrangian leading to Eqs.(10) and (11) is now expressed as

$$L_{\text{eff}} = \frac{1}{2} \sum_{j=1}^2 \left[ \dot{\mathbf{R}}_j^2 - R_j^2 + \dot{w}_j^2 - w_j^2 - \frac{1}{w_j^2} \left( 1 + \frac{g}{2\pi} \right) \right] - \frac{\Lambda g}{\pi(w_1^2 + w_2^2)} e^{-\frac{(\mathbf{R}_1 - \mathbf{R}_2)^2}{w_1^2 + w_2^2}}. \quad (12)$$

Let us define the center-of-mass of two components  $\mathbf{R}_0 = \frac{\mathbf{R}_1 + \mathbf{R}_2}{\sqrt{2}}$  and the relative displacement  $\mathbf{Q} = \frac{\mathbf{R}_2 - \mathbf{R}_1}{\sqrt{2}}$ , and suppress the global translational degree of freedom ( $\mathbf{R}_0 = \dot{\mathbf{R}}_0 = 0$ ). Owing to rotational symmetry the angular momentum  $M$  is a constant of motion:  $M = \frac{\partial L_{\text{eff}}}{\partial \dot{\theta}} = Q^2 \dot{\theta}$  with  $\theta$  as a polar angle for  $\mathbf{Q}$ . Choosing a synchronous width dynamics ( $w_1 = w_2 = w$ ) with the canonical momentum  $p_w = \frac{\partial}{\partial \dot{w}} (L_{\text{eff}} - M \dot{\theta})$ , the effective Hamiltonian is given by

$$H_{\text{eff}} = \frac{1}{2} P_Q^2 + \frac{1}{4} p_w^2 + V(Q, w)$$

with

$$V(Q, w) = \frac{1}{2} Q^2 + \frac{M^2}{2Q^2} + w^2 + \frac{1}{w^2} \left( 1 + \frac{g}{2\pi} \right) + \frac{\Lambda g}{2\pi w^2} e^{-\frac{Q^2}{w^2}}. \quad (13)$$

Equation (13) indicates: In the absence of ICI ( $\Lambda = 0$ ),  $w$  and  $Q$  show independent regular oscillations around the minima of individual potentials,  $w = w_0 = (1 + g/2\pi)^{1/4}$  and  $Q = Q_0 = \sqrt{M}$ , respectively. Interestingly, however, for any set of values  $M$  and  $g$  the typical oscillation frequencies  $\omega_Q^0$  for  $Q$  and  $\omega_w^0$  for  $w$  are always degenerate:  $\omega_Q^0 = \omega_w^0 = 2$  (in unit of frequency of the harmonic potential,  $\omega$ ). Therefore, by switching on a small but non-vanishing ICI, dynamics of  $w$  and that of

$Q$  will be strongly mixed to remove the degeneracy and normal modes, fat (optical) and skinny (acoustic), are formed. In the fat mode,  $Q$  and  $w$  change in anti-phase, namely the decrease of the relative displacement is accompanied by the blowup of WPs, while in the skinny mode,  $Q$  and  $w$  change in phase (see Fig.1(b)). The emergence of the fat mode brings about the wave interference between adjacent WPs as shown in Fig.1(a). This provides a mechanism leading to the breakdown of Pérez-García *et al.*'s picture of interacting particles.




FIG. 2: Upper panel: Time evolution of WP width ( $w_j$ ).  $g = 2\pi$ ,  $\Lambda g = 3$  and angular momentum  $M = 9$ .  $\Omega$  is a tunable parameter. (a)  $\Omega = 0$  (dotted line) and  $\Omega = 100$  (solid line), (b)  $\Omega = 3$ , (c)  $\Omega = 10$ ; Lower panel: Time evolution (up to  $t = 100[\omega^{-1}]$ ) of initial three Gaussian WPs in 3-component BEC with  $g = 2\pi$ ,  $\Lambda g = 3$ ,  $\Omega = 100$  and  $M = 9$ . See the persistence of a particle picture.

To suppress the wave interference between adjacent WPs and recover the particle picture, we proceed to introduce a time-periodic ICI with zero time-average and the amplitude  $\Lambda g$ ,

$$g_{ij} = \Lambda g \cos(\Omega t) \quad (i \neq j) \quad (14)$$

rather than sticking to the static ICI. Here the frequency scale  $\omega_w^0 (= 2)$  is essential. If  $\Omega \gg \omega_w^0$ , the width will feel actually no ICI, showing a stable and regular breathing proper to a single component BEC with a harmonic trap. For  $\Omega$  near but larger than  $\omega_w^0$ , one can expect a sufficiently long stable breathing until the time ( $\gg 2\pi/\omega_w^0$ ) that a blowup of WP will start. For  $\Omega$  equal or less than  $\omega_w^0$ , the breathing oscillation will quickly blow up leading to the wave interference.

To verify the above conjecture, we have numerically integrated 3-component NSE (GP equation) with a harmonic trap in two dimensions in (1) with use of split-step Crank-Nicholson method [12]. Initial data are coordinates  $X_j = R \cos(2\pi(j-1)/3)$ ,  $Y_j = R \sin(2\pi(j-1)/3)$ .




FIG. 3: Time evolution of WP width ( $w_j$ ) (upper panel) and corresponding power spectrum (lower panel).  $g = 2\pi$ ,  $\Omega = 5$  and  $M = 9$ .  $\Lambda g$  is a tunable parameter. (a)(a')  $\Lambda g = 0$ , (b)(b')  $\Lambda g = 2$ , (c)(c')  $\Lambda g = 4$ .

1)/3) and velocities  $V_{jX} = -R\omega \sin(2\pi(j-1)/3)$ ,  $V_{jY} = R\omega \cos(2\pi(j-1)/3)$  with  $R^2\omega = M/3$  ( $M$ : angular momentum) and  $j = 1, 2, 3$ . In short these are placed on the vertices of an equilateral triangle. We have also calculated expectation values of the widths using  $w_{jX}^2 = \int ((x - \langle x \rangle)^2 |\psi_j|^2) d\mathbf{r}$ ,  $w_{jY}^2 = \int ((y - \langle y \rangle)^2 |\psi_j|^2) d\mathbf{r}$  for different times. In Figs. 2 and 3 the mean value  $w_j \equiv (w_{jX} + w_{jY})/2$  is plotted as a function of time. Note: even for a single-component NSE[6, 7], the initial circular symmetry of WP is slightly broken during the time evolution, due to the nonlinear *intra-component* interaction; Each of the expectations  $w_{jX}$  and  $w_{jY}$  is described by a superposition of modes with nearly-equal frequencies  $\omega_w^0$  and  $\sqrt{(1 + g/4\pi)/(1 + g/2\pi)}\omega_w^0$  and shows a small beating, which however can be suppressed by choosing the above mean value  $w_j$ .

Figure 2(a) shows a blowup of the breathing oscillation for the static ICI ( $\Omega = 0$ ) and a recovery of stable oscillation for  $\Omega \gg \omega_w^0$ , and Figs. 2(b) and 2(c) show oscillations with no blowup for  $\Omega \gtrsim \omega_w^0$ . For a fixed value of  $\Omega$  in the range  $\Omega \simeq \omega_w^0$ , we see a transition from regular to chaotic breathing oscillations when the amplitude  $\Lambda g$  is increased. In fact, in the case of  $\Omega = 5$ , the oscillation is irregular for  $\Lambda g = 2$  and 4 (see Figs. 3(b) and 3(c)), while it is regular for  $\Lambda g = 0$  (see Fig. 3(a)). The corresponding power spectra show a transition from a distinguished line structure in Fig. 3(a') to broad ones in Fig. 3(b') and




FIG. 4: Phase diagram in  $\Lambda g$ - $\Omega$  space. ( $g = 2\pi$  and  $M = 9$ )  $\times$ ,  $\circ$ ,  $\square$  stands for blowup, chaotic and regular regions, respectively. (Data are increased in the left half of the diagram for use of detailed discussions.)

3(c').

We have systematically investigated the wave dynamics starting commonly from the equilateral triangle configuration as above. Figure 4 is a phase diagram in  $\Lambda g$ - $\Omega$  space, which shows three distinct regions, (i) blowup, (ii) regular without blowup and (iii) chaotic without blowup. We find the decrease of the blowup region as  $\Omega$  is increased beyond  $\omega_w^0$  and the emergence of chaos without the blowup as  $\Lambda g$  is increased. Contrary to our simple expectations, the phase diagram involves rich information to be explored: The regular and chaotic oscillations without blowup are observed even for  $\Omega < \omega_w^0$ , and the blowup occurs at resonances,  $\Omega = \omega_w^0$  and  $2\omega_w^0$ , even for a small amplitude  $\Lambda g$ . However, Fig. 4 clearly conveys that the three-component BEC with the oscillating ICI restores a picture of interacting particles and can exhibit the transition from regular to chaotic motions without the blowup. Thus we have obtained a model of conservative chaos constructed from the three-component BEC with a harmonic trap in 2 dimensions.

To develop Pérez-García *et al.*'s idea, we have examined the multi-component repulsive BEC in a 2-d harmonic trap. In the absence of the inter-component interaction (ICI), the wavepacket (WP) breathing and the motion of relative distance between WPs have oscillation frequencies always degenerate for any set of the angular momentum and intra-component interaction. The non-zero ICI removes the degeneracy and induces the fat mode that breaks a picture of soliton molecules. We have therefore proposed a new model with a time-periodic ICI with zero time-average, which can elongate the time interval without any blowup of WPs. Provided that the frequency is near the characteristic breathing frequency ( $\omega_w^0$ ), a transition from regular to chaotic oscillations occurs as the amplitude of ICI is increased. For attractive BECs *without a trap* or in free space, the oscillating nonlinearity *with non-zero average* is known to stabilize high-dimensional WPs[13, 14]. In a similar way, in the case of the multi-component repulsive BEC *with a trap*,

the analogous picture holds for high-dimensional WPs mutually interacting through time-periodic ICI *with zero average*. The amplitude of ICI is found to control the chaoticity of breathing motions of WPs. The analysis of chaos in space coordinates and the comparison of the

present results based on NSE with the effective dynamics with a few degrees-of-freedom in Eqs. (10) and (11) will constitute subjects which we intend to study in future.

K.N. is grateful to M. Lakshmanan for valuable conversations in the early stage of the present work.

- 
- [1] M.R.Andrews, C.G.Townsend, H.J.Miesner, D.S.Durfee, D.M.Kurn and W.Ketterle, *Science* **275**, 637(1997)
  - [2] M.R.Andrews, D.M.Kurn, H.-J.Miesner, D.S.Durfee, C.G.Townsend, S.Inouye, and W.Ketterle, *Phys. Rev. Lett.* **79**, 553(1997); **80**, 2967(1998)
  - [3] D.S.Hall,M.R.Matthews,J.R.Ensher,C.E.Wieman and E.A.Cornell, *Phys. Rev. Lett.* **81**, 1539(1998)
  - [4] K.E.Strecker,G.B.Partridge,A.G.Truscott and R.G.Hulet, *Nature(London)* **417**, 150(2002)
  - [5] L.Khaykovich, F.Schreck,G.Ferrari,T.Bourdel,J.Cubizolles, L.D.Carr,Y.Castin and C.Salomon, *Science* **296**, 1290 (2002)
  - [6] V.M.Pérez-García, J.I.Cirac, M.Lewenstein and P.Zoller, *Phys. Rev. Lett.* **77**, 27(1996)
  - [7] V.M.Pérez-García, H.Michinel, J.I.Cirac, M.Lewenstein and P.Zoller, *Phys. Rev. A* **56**, 1424(1997)
  - [8] J.J. García-Ripoll, V.M.Pérez-García and V.Vekslerchik, *Phys. Rev. E* **64**, 056602(2001)
  - [9] V.M.Pérez-García and V.Vekslerchik, *Phys.Rev.E* **67**, 061804(2003)
  - [10] G.D.Montesinos,V.M.Pérez-García and H.Michinel, *Phys. Rev. Lett.* **92**, 133901-1(2004)
  - [11] V.M.Pérez-García, *Physica D* **191**,211(2004)
  - [12] See, e.g., P.Muruganandam and S.K.Adhikari, *J. Phys. B:At. Mol. Opt. Phys.* **36**, 2501(2003)
  - [13] H.Saito and M.Ueda, *Phys. Rev. Lett* **90**, 040403(2003)
  - [14] F.K.Abdullaev, J.G.Caputo, R.A.Kraenkel and B.A.Malomed, *Phys. Rev. A* **67**, 013605(2003)